

UC Riverside

UC Riverside Previously Published Works

Title

GnRH Receptor Expression and Reproductive Function Depend on JUN in GnRH Receptor-Expressing Cells.

Permalink

<https://escholarship.org/uc/item/49f3q50q>

Journal

Endocrinology, 159(3)

ISSN

0013-7227

Authors

Jonak, Carrie R
Lainez, Nancy M
Boehm, Ulrich
et al.

Publication Date

2018-03-01

DOI

10.1210/en.2017-00844

Peer reviewed

GnRH receptor expression and reproductive function depend on JUN in the GnRH receptor-expressing cells

Carrie R. Jonak¹, Nancy M. Lainez¹, Ulrich Boehm², and Djurdjica Coss¹

¹Division of Biomedical Sciences; School of Medicine, University of California, Riverside; Riverside, CA 92521. ²Experimental Pharmacology, Center for Molecular Signaling (PZMS), Saarland University School of Medicine, 66421 Homburg, Germany.

Abbreviated title: Gonadotrope c-Jun regulates reproduction

Corresponding author: Djurdjica Coss

Division of Biomedical Sciences,
School of Medicine,
303 SOM Research Building
University of California, Riverside;
Riverside, CA 92521
Tel: 951 827-7791, Fax: 951 827-2477, E-mail: djurdjica.coss@ucr.edu

Key words: c-Jun, reproduction, conditional knockout, gonadotrope, AP-1, GNRH, GnRH receptor,

Disclosure Summary:

The authors declare no competing financial interests and have nothing to disclose.

Acknowledgements:

This research was supported by NIH grant R01 HD057549 and UCR School of Medicine start-up funds (to DC). University of Virginia Ligand Core is supported by U54 HD28934. The authors thank Dr. Randy Johnson for the cJun^{flox/flox} mice, Dr. Greg Anderson for GnRH antibody, and Dr. Pamela Mellon for the LβT2 cells. We are grateful to Dr. Ameae Walker for thoughtful discussion.

Abstract

Gonadotropin-releasing hormone (GnRH) from the hypothalamus regulates synthesis and secretion of luteinizing hormone (LH) and follicle-stimulating hormone (FSH) from the anterior pituitary gonadotropes. LH and FSH are heterodimers, comprised of a common α -subunit and unique β -subunits, which provide biological specificity and are limiting components of the mature hormone synthesis. Gonadotrope cells respond to GnRH via specific expression of the GnRH receptor. GnRH induces the expression of gonadotropin genes and of the GnRH receptor by activation of specific transcription factors. The JUN (c-Jun) transcription factor binds to AP-1 sites in the promoters of target genes and mediates induction of the FSH β gene and of the GnRH receptor in the gonadotrope-derived cell lines. To analyze the role of JUN in reproductive function *in vivo*, we generated a new mouse model that lacks JUN specifically in GnRH receptor-expressing cells (JUN-cKO). JUN-cKO mice displayed profound reproductive anomalies such as reduced LH levels resulting in lower gonadal steroid levels, longer estrous cycles in females, and diminished sperm numbers in males. Unexpectedly, FSH levels were unchanged in these animals, while GnRH receptor expression in the pituitary was reduced. Steroidogenic enzyme expression was reduced in the gonads of JUN-cKO mice, likely as a consequence of reduced LH levels. GnRH receptor driven Cre activity was detected in the hypothalamus, but not in GnRH neuron. Female, but not male, JUN-cKO mice exhibited reduced GnRH expression. Taken together, our results demonstrate that GnRH receptor expression levels depend on JUN and are critical for reproductive function.

Precis

Knock-down of JUN in the GnRH receptor-expressing cells leads to diminished reproductive capacity, reduced GnRH receptor expression and lower serum LH in male and female mice.

Introduction

Mammalian reproduction is regulated by the hypothalamic-pituitary-gonadal (HPG) axis. The hypothalamic decapeptide gonadotropin-releasing hormone (GnRH) is the final brain output that regulates both expression and secretion of gonadotropins, luteinizing hormone (LH) and follicle stimulating hormone (FSH) from the anterior pituitary gonadotropes (1). This function is mediated by gonadotrope-specific expression of the GnRH receptor, which belongs to the rhodopsin family of seven transmembrane G-protein coupled receptors (2). LH and FSH in turn stimulate steroidogenesis and gametogenesis in the gonads (3, 4).

Gonadotropin levels are primarily regulated by transcription of their unique β -subunits, which provide biological specificity. Alternations in the transcription of β -subunits correlate with changes in the concentration of the mature hormones in the circulation (5, 6). The β -subunits heterodimerize with a common α -subunit to form the mature glycoproteins (7). GnRH induces LH β (*Lhb*), FSH β (*Fshb*) and GnRH receptor (*Gnrhr*) transcription via induction of specific immediate-early genes: EGR1 that regulates *Lhb* transcription; and FOS and JUN, which activate both *Fshb* and *Gnrhr* transcription (4). The FOS and JUN transcription factors form the AP-1 heterodimer, which is rapidly and transiently activated (8). Both mouse and human *Fshb* and *Gnrhr* genes are induced by GnRH via AP-1 (9-13). Transcriptome analysis demonstrated that

AP-1 members are strongly induced by GnRH in L β T2 cells (14) and in primary rat gonadotrope cells (15).

Responsiveness of the *Fshb* gene to GnRH is conveyed by AP-1 response elements in the proximal promoter (9,16-19). GnRH induces FOS (c-Fos), FOSB, JUN (c-Jun) and JUNB, but not JUND in the L β T2 cell line, a model of mature gonadotropes. A combination of these factors binds the AP-1 site in the *Fshb* promoter (9). In the α T3 gonadotrope cell line, GnRH regulates *Gnrhr* expression via AP-1, as well (11, 20). JUN homodimer, or a heterodimer with FOS, FOSB, FRA1 or FRA2, binds the mouse *Gnrhr* promoter at two different sites (13, 21). AP-1 heterodimer of JUN and FOS also regulates expression of the human *GNRHR* gene by GnRH (22).

Although gonadotrope cell models, such as L β T2 and α T3 cells, facilitated identification of transcription factors that lead to induction of gonadotrope genes, it is critical to determine the roles of these transcription factors *in vivo*. LH β induction by GnRH is mediated by the EGR1 transcription factor. EGR1 is an immediate early gene and a member of the zinc finger family of transcription factors. EGR1 plays a non-redundant role in reproduction, and other family members are unable to compensate. Consistent with this, global EGR1 knockout mice are infertile and lack LH expression resulting in blunted sex steroid hormone synthesis (23, 24). FOS also plays non-redundant roles in reproduction *in vivo* (25). In the pituitary, FOS is critical for gonadotropin gene expression, while expression of another glyco hormone subunit, TSH β (*Tshb*) is not affected. In the hypothalamus, FOS is expressed in both kisspeptin and GnRH neurons during the preovulatory surge and can be used as a marker of their activation (26-28). FOS is necessary for normal kisspeptin neuron numbers and *Kiss1* expression, primarily in the female, while GnRH neuron location, axon targeting or gene expression do not depend on FOS (25).

Since JUN is an obligatory heterodimerization partner of FOS for DNA binding (8), we used c-Jun^{flox/flox} mice crossed to GnRH receptor Cre animals to create mice that lack JUN specifically in the GnRH receptor-expressing cells. These conditional knockout mice, JUN-cKO, were used to analyze the reproductive physiology and determine the cell-specific role of JUN in reproduction.

Materials and Methods

Cell lines and transient transfection

LβT2, a gift from Dr. Pamela Mellon (UCSD), were maintained in DMEM with 10% FBS at 37°C and 5% CO₂. The line was authenticated with RT-PCR based expression analysis of endogenous gonadotropin β subunits. For transfection, LβT2 cells were plated in 12-well plates one day prior to transfection with FuGENE 6 transfection reagent (Roche Molecular Biochemicals, Indianapolis, IN), 0.1 μg expression vectors, 0.5 μg of a luciferase-reporter plasmid (reported previously (17, 29-31)) and 0.1 μg of TK β-galactosidase, a reporter plasmid driven by a Herpes virus thymidine kinase (TK) promoter as a control for transfection efficiency. cDNA for the AP-1 transcription factors were in the same backbone under the same promoter, and their expression was evaluated by western blot. Forty-eight hours after transfection, cells were lysed with 100 nM KPO₄ buffer containing 0.2% Triton X-100 luciferase activity measured on a luminometer (Veritas Microplate luminometer from Turner Biosystems) by injecting 100 μl of buffer containing 25 mM Tris pH 7.8, 15 mM MgSO₄, 10 mM ATP, and 65 μM luciferin into each well. Using the Tropix Galactolight β-galactosidase assay (Applied Biosystems, Foster City, CA) and following the manufacturer's instructions, β-galactosidase activity was measured subsequently. Transfections

were performed in triplicate and repeated a minimum of three times. 1-way ANOVA statistical analysis with Tukey's posthoc test was performed using the JMP program with significance set at $p < 0.05$.

Animals

Mice lacking c-Jun in GnRH receptor-expressing cells were obtained by crossing c-Jun^{flox/flox} mice with GnRH-Receptor-Cre (GRIC) mice. Briefly, c-Jun^{flox/flox} mice, in which the only coding exon of the *c-Jun* allele is flanked by *LoxP* sites (32, 33), were created by Dr. Randall Johnson (UCSD, California). Gnhr^{tm1(cre)Uboe} mice (GnRH receptor-internal ribosome entry site-Cre, GRIC) carry a knock-in GnRH receptor allele fused to an internal ribosome entry site and a Cre transgene. GRIC drives Cre expression in pituitary gonadotrope cells (34). Since some Cre expression is also observed in male germ cells in these animals (35), the GRIC allele was always introduced via the female. Homozygous c-Jun^{flox/flox} Cre⁺ mice served as experimental mice, while Cre⁻ littermates were used as controls. TdTomato reporter mice, *Gt(ROSA)26Sor^{tm9(CAG-tdTomato)Hze}/J*, were obtained from Jackson laboratory (strain 007909) and crossed to GRIC mice to analyze Cre activity. Animals were maintained under a 12-hour light, 12-hour dark cycle and received food and water *ad libitum*. All experiments were performed with approval from the University of California Animal Care and Use Committee and in accordance with the National Institutes of Health Animal Care and Use Guidelines using 8-week-old animals, unless indicated otherwise. Males and females were analyzed separately to determine potential sex differences. At least 6 animals per sex per genotype were analyzed and statistical differences between Cre⁺ and Cre⁻ were determined by Student's T-test and Tukey's test for multiple comparison.

Fertility studies – 8-week-old Cre⁺ and Cre⁻ male or female mice were individually paired with an adult C57BL/6 mouse of the opposite sex, and the presence of litters was monitored daily over a period of 4 months. Additionally, starting at 8 weeks of age, a separate cohort of female mice was assessed for estrous cycle stage with daily vaginal smears for 5 weeks.

Sperm count - The epididymides were dissected, macerated, incubated in 1 ml DMEM at room temperature for 30 minutes with shaking. Sperm was cleared with a 70 µm cell strainer, diluted with sterile water and counted with a haemocytometer.

Histological analyses and immunohistochemistry

Ovaries and testes were fixed overnight at 4°C in 4% paraformaldehyde or Bouin's fixative, respectively. Tissues were dehydrated in ethanol, embedded in paraffin, cut into 10 µm thick sections, floated onto UltraClear™ Plus Microslides (Denville Scientific Inc, Holliston, Massachusetts) and stained with hematoxylin and eosin.

Pituitaries were fixed in 4% paraformaldehyde, embedded in paraffin, and cut to 10 µm. Slides were deparaffinized in xylene and rehydrated. Antigen unmasking was performed by heating for 10 minutes in a Tris-EDTA-0.3% Triton X and endogenous peroxidase was quenched by incubating for 10 minutes in 0.3% hydrogen peroxide. Slides were then blocked with 20% goat serum and incubated with primary antiserum against LH (1:300 raised in rabbit, National Hormone and Peptide Program, NIDDK) overnight at 4°C. After PBS washes, slides were incubated with biotinylated goat anti-rabbit IgG (1:300, BA-1000, Vector Laboratories, Burlingame, CA) for 30 minutes. The Vectastain ABC elite kit (Vector Laboratories) was used per manufacturer's instructions, after which the DAB peroxidase kit was used for colorimetric staining. Slides were dehydrated in ethanol and xylene, and cover-slipped with Vectamount (Vector Laboratories).

To visualize costaining of TdTomato and pituitary hormones, pituitaries were fixed in 4% paraformaldehyde, frozen in OCT, and cut to 12 μ m sections using Leica cryostat. Hypothalami were sectioned to 30 μ m sections for GnRH staining. Slides were blocked with 20% goat serum and incubated with primary antibodies against LH or FSH (1:300 raised in rabbit, National Hormone and Peptide Program, NIDDK) or GnRH (provided kindly by Greg Anderson, University of Otago; Dunedin, New Zealand (36)) overnight at 4°C. After PBS washes, slides were incubated with biotinylated goat anti-rabbit IgG (1:300, BA-1000, Vector Laboratories, Burlingame, CA) for 30 minutes; followed by streptavidin-Cy5 (1:500, Molecular Probes, Thermo Fisher) for 30 minutes. Secondary antibody-only controls were performed and determined that endogenous TdTomato expression was strong for visualization and that its emission in the TdTomato/Rhodamine channel overlaps with FITC/Alexa 488 channel. Thus, Streptavidin-Cy5 was used for visualization of LH-, FSH- or GnRH-expressing cells and slides cover-slipped using Vectasheild (Vector Laboratories). To determine percent co-expression, we counted how many of the hundred LH- or FSH-containing cells express TdTomato and vice versa. We counted at least 3 non-overlapping fields of view in 3 different sections per mouse (=9 fields), and stained pituitaries from 3 male and 3 female Cre+ mice.

qPCR analyses

Tissues were dissected, total RNA extracted and reverse transcribed using Superscript III (Invitrogen, CA). qPCR was performed using an iQ SYBR Green supermix and an IQ5 real-time PCR machine (Bio-Rad Laboratories, Hercules, CA) with primers listed in Table 1 under the following conditions: 95°C for 15 min, followed by 40 cycles at 95°C for 20 sec, 56°C for 30 sec, and 72°C for 30 sec. A standard curve with dilutions of 10 pg/well, 1 pg/well, 100 fg/well, and 10

fg/well of a plasmid containing LH β , or FSH β cDNA, was generated in each run with the samples. The amount of the gene of interest was calculated by comparing the threshold cycle obtained for each sample with the standard curve generated in the same run. Replicates were averaged and divided by the mean value of the GAPDH housekeeping gene in the same sample using $\Delta\Delta C_t$ method. After each run, a melting curve analysis was performed to confirm that a single amplicon was generated in each reaction. Statistical differences in expression between genotypes were determined by Student's T-test, and Tukey's HSD for multiple comparisons using JMP software (SAS Institute; Cary, North Carolina).

Hormone analyses

For serum collection, mice were sacrificed between 9-11 am by isoflurane inhalation and blood was obtained from the inferior *vena cava*. The blood was left to coagulate for 15 minutes at room temperature, and then centrifuged at 2000 RCF for 15 minutes for serum separation. Hormone assays were performed by the University of Virginia, Ligand Core. The University of Virginia Center for Research in Reproduction Ligand Assay and Analysis Core is a fee-for-service core facility and is in part supported by the Eunice Kennedy Shriver NICHD/NIH Grant U54-HD28934. LH was analyzed using a sensitive two-site sandwich immunoassay (37), and mouse LH reference prep (AFP5306A; provided by Dr. A.F. Parlow and the National Hormone and Peptide program) was used as standard. FSH was assayed by RIA using reagents provided by Dr. A.F. Parlow and the National Hormone and Peptide Program, as previously described (38). Mouse FSH reference prep AFP5308D was used for assay standards. Steroid hormone levels were analyzed using validated commercially available assays, information for which can be found on the core's website: <http://www.medicine.virginia.edu/research/institutes-and-programs/crr/lab->

facilities/assay-methods-page and reported in (39). Limits of detection were 0.24 ng/ml for LH, 2.4 ng/ml for FSH, 3 pg/ml for estradiol, and 10 ng/dL for testosterone. Intra- and inter-assay coefficients of variation were 6.4%/8.0%, 6.9%/7.5%, 6.0%/11.4% and 4.4%/6.4% for the LH, FSH, estrogen (E2) and testosterone (T), respectively. For the assays used for this manuscript, inter-assay coefficients of variation data are the result of 30 assays for LH and FSH, and 60 assays for E2 and T. Six animals per group were used for each hormone analysis. Statistical differences in hormone levels between wild-type and null group were determined by Student's T-test, and Tukey-Kramer HSD for multiple comparisons using JMP software (SAS Institute; Cary, North Carolina).

Results

JUN induces FSH β and GnRH receptor reporters in L β T2 gonadotrope cell line.

Given that the AP1 family of transcription factors is comprised of four FOS members (FOS (c-Fos), FOSB, FRA1 and FRA2) and three JUN members (JUN (c-Jun), JUNB and JUND), combinatorial heterodimerization of these provides a variety of different factors that can induce target genes. GnRH induces all family members in gonadotropes, except for JUND (9). Since AP-1 heterodimers bind FSH β (*Fshb*) and GnRH receptor (*Gnrhr*) promoters using EMSA (9, 13, 21), we first analyzed the level of induction of these target genes in gonadotropes with different combinations of AP-1 factors. cDNAs for the AP-1 transcription factors were cloned in the same vector backbone under the same promoter, and their expression was confirmed by western blot (data not shown). We also compared the induction with AP-1 overexpression to the induction by GnRH (G, Fig. 1). Since GnRH receptor reporter induction by GnRH was previously analyzed

using α T3-1 cells, a model of immature gonadotrope, we determine the level of induction in the model of mature gonadotrope, L β T2 cells. GnRH induced FSH β reporter 6.2 fold, and GnRH receptor reporter 2.4 fold. FRA1 (F1) and FRA2 (F2) did not induce FSH β (Fig. 1A) or GnRH receptor (Fig. 1B) expression either alone, nor more highly in combination with either JUN or JUNB compared to JUN or JUNB alone. JUN in combination with FOS or FOSB induced FSH β reporter to similar levels compared to the induction observed with GnRH treatment (Fig. 1A). JUN heterodimers induced FSH β to higher levels compared to JUNB heterodimers with FOS or FOSB. GnRH receptor, on the other hand, was induced to similar levels by either JUN or JUNB heterodimers with FOS or FOSB (Fig. 1B). In L β T2 cells, GnRH receptor is induced by GnRH 2.4 fold, a similar level observed with AP-1 overexpression. Since JUN induces both AP-1 gene targets in gonadotrope-derived cell line, we next crossed c-JUN^{flox/flox} mice to GnRH receptor Cre (GRIC) animals to analyze the role of JUN in gonadotropes *in vivo*.

Reduced reproductive capacity but normal gonadotrope numbers in mice lacking JUN in GnRH receptor-expressing cells. Previous studies successfully used the GRIC allele to express Cre recombinase in gonadotropes to analyze transcription factors' roles in gonadotropin gene expression (40-42). We used the GRIC allele to knockdown JUN and create a conditional JUN knockout (JUN-cKO). Because JUN is an immediate early gene that is expressed at a very low basal level, undetectable by immunostaining, we were unable to reliably demonstrate JUN knockdown in the gonadotrope. Thus, to analyze Cre activity in the gonadotrope, and co-expression of Cre and gonadotropin hormones, we used TdTomato reporter mice in which TdTomato is specifically induced in Cre-expressing cells, following Cre-mediated excision of the stop codon. Immunohistochemistry of frozen pituitary sections with antibodies to gonadotropin

hormones, revealed faithful expression of the TdTomato fluorescence; that 98% of TdTomato expressing cells also express LH or FSH, consistent with the previous report (34). Furthermore, 88% of cells that contain LH express TdTomato (Fig. 2A, white arrowheads indicate LH-containing cells lacking TdTomato expression). 76% of FSH-containing cells express TdTomato (Fig. 2B).

JUN-cKO animals exhibited profound changes in their reproductive physiology; females had significantly longer estrous cycles, 7.4 days, compared to 4.4 days per cycle in controls (Fig. 3A, representative females' stage of the estrous cycle over a 33-day period; Fig. 3B, days per cycle in 6 females per genotype; JUN-cKO, cKO; control, Ctr). Male JUN-cKO mice had a 43% lower sperm count compared to controls (Fig. 3C). JUN-cKO mice also displayed longer time intervals in between litters, when paired with wild type C57BL/6 mice of the opposite sex (Fig. 3D, female cKO data presented, male data not shown).

To assess the role of JUN in gonadotrope differentiation, we stained pituitaries from JUN-cKO and control mice for LH to determine the number of gonadotropes. The morphology and size of JUN-cKO (cKO, Cre+) and control (Ctr, Cre-) pituitaries were indistinguishable (Fig. 4A). We then counted gonadotropes and determined that animals of both sexes and both genotypes contained the same numbers of gonadotropes (Fig. 4B). Therefore, the lack of JUN in the gonadotropes did not affect gonadotrope numbers. The JUN-cKO and control animals were of the same size and weight (data not shown). Therefore, despite the same number of gonadotropes, the lack of JUN in gonadotrope cells results in subfertility in both sexes.

JUN-cKO mice have reduced LH levels. Analyses of gonadotropin levels in the circulation revealed that JUN-cKO males exhibited 49% lower serum LH compared to control males, while

the LH concentration in JUN-cKO diestrus females was reduced by 56% compared to control females in diestrus (Fig 5A). Although GnRH induces FSH β via FOS and JUN in the gonadotrope-derived cell line, FSH levels were the same in both JUN-cKO and control males and in JUN-cKO and control diestrus females (Fig 5B). Steroid hormone levels were reduced; testosterone was lower in males, while estradiol was lower in females (Fig. 5C), likely due to reduced LH levels in the circulation.

We also analyzed gonadotrope gene expression at 8 weeks of age. Consistent with the reduction in LH concentration in the circulation, *Lhb* mRNA level was 29% lower in JUN-cKO males (Fig. 6A) and 62% lower in JUN-cKO diestrus females compared to Cre- littermate controls (Fig. 6B). Consistent with unaltered FSH levels, there was no difference in *Fshb* expression between genotypes (Fig. 6C, D). Expression of the *Gnrhr* (GnRH receptor) mRNA, however, was reduced by 28% in JUN-cKO males (Fig. 6E) and by 56% in JUN-cKO females (Fig. 6F). Expression of the common *Cga* subunit (α GSU) that heterodimerizes with both LH β and FSH β was unaffected (Fig. 6G, H). Previous studies analyzing *Lhb* expression did not reveal a role for the FOS and JUN AP-1 family, while the importance of AP-1 in GnRH receptor induction is well established (13,43). Our results may point to a role for AP-1 in *Lhb* expression. On the other hand, concomitant reduction of both *Lhb* and *Gnrhr* expression in both JUN-cKO males and females may implicate diminished GnRH receptor levels in lower *Lhb* mRNA. This is consistent with previous studies postulating that the receptor concentration correlates with LH β levels (44).

Reduced LH target genes in the gonads of JUN-cKO mice. We next analyzed potential downstream effects of reduced LH levels in the gonads in both males and females at 12 weeks of age. Male JUN-cKO mice had a 22% reduction in seminal vesicle weight compared to controls

(Fig. 7A), which is consistent with reduced intratesticular testosterone levels (Fig. 7B). We also examined the expression of steroidogenic enzymes, which are induced by LH signaling. While *Star* (Steroidogenic acute regulatory protein, StAR) expression was unchanged, expression of *Cyp11* and *Cyp17* was reduced by 20% and 25%, respectively (Fig 7C, D, E). Expression of the FSH target gene in the testis, *Shbg* (Sex hormone binding globulin; androgen-binding protein, ABP) was unaffected, consistent with the unperturbed FSH levels in the circulation (Fig. 7F). We observed lower sperm numbers, as shown above. Testosterone levels, that were reduced due to the reduction in LH concentration, are necessary for spermatogenesis and for the maintenance of the blood-testis barrier. Blood-testis barrier is established via expression of tight junction proteins from the Claudin family (45,46). Expression of claudin 11 (*Cldn11*) did not change (data not shown). Claudin 3 (*Cldn3*) expression is regulated by androgens (47), however despite a decrease in testosterone, expression of *Cldn3* was not significantly reduced (Fig. 7G, p=0.1). Given that sperm numbers in the epididymides were diminished, we assessed markers for several stages of spermatogenesis (48) and determined that the early stage spermatogenesis marker *Sycp3* to be unchanged, while later stage markers such as *Spert* and *Elp* were reduced in JUN-cKO males by 31% and 36% compared to the controls, respectively (Fig. 7H, I, J). Histological analyses of the testes uncovered small number of abnormal seminiferous tubules (~5%) lacking mature sperm in the JUN-cKO males (Fig. 7L). Thus, lack of JUN in GnRH receptor-expressing cells in JUN-cKO males causes lower expression of steroidogenic enzymes and reduced levels of the late stage spermatogenesis markers, corresponding to reduced sperm count.

The ovaries of JUN-cKO mice weighed 37% less than control ovaries and contained fewer corpora lutea (Fig. 8A, B, C). JUN-cKO females expressed 43% lower level of the LH target gene *Cyp17a1* (Fig. 8D), while the FSH target gene *Cyp19a1* (aromatase) was unchanged in the ovaries

(Fig. 8E). Given that antral stage of folliculogenesis is not affected corresponding to unaltered FSH, fewer corpora lutea may stem from reduction in prolactin levels, since prolactin is necessary for corpus luteum function in rodents (49, 50). We measured expression of prolactin (*Prl*) in the pituitary and determined that *Prl* mRNA is reduced by 67% in JUN-cKO female mice (Fig. 8F). Therefore, female as well as male gonads from JUN-cKO animals exhibit a phenotype corresponding to diminished reproductive capacity.

Cre activity in the hypothalamus. The lack of an effect on *Fshb* expression and FSH levels in the circulation was unexpected, given previous evidence in the literature. In addition to inadequate Cre activity in a portion of gonadotrope cells, other JUN family members such as JUNB, that is also induced by GnRH (9), may compensate for the loss of JUN. Although JUN and JUNB exert non-overlapping functions in other tissues as evidenced by the different phenotypes of the respective knockout mice (32, 51), they may be able to substitute for each other in this scenario. To assess a possible compensatory increase in JUNB expression, we analyzed the level of *Junb* mRNA in the pituitaries of JUN-cKO and of control males and females. In both sexes, JUN-cKO animals exhibited an increase in JUNB expression in the pituitary (Fig. 9). Therefore, JUNB increase may be able to compensate for the loss of JUN for *Fshb* but not for *Gnrhr* expression.

On the other hand, reduced expression of *Gnrhr* and *Lhb* may stem from extrapituitary sites. GnRH receptor is expressed in several hypothalamic nuclei and may be expressed in GnRH neurons themselves (52-58). We used TdTomato reporter mice to determine activity of Cre recombinase in the hypothalamus. We also performed immunostaining for GnRH to detect GnRH neurons and determine whether TdTomato is expressed in GnRH neurons following Cre excision of the stop codon. Coronal sections of the mediobasal hypothalamus demonstrated that TdTomato

was expressed in the arcuate nucleus in GRIC+ animals, while GnRH axon terminals were located in the median eminence (Fig 10A, GnRH, green; TdTomato, red). Staining of the preoptic area detected GnRH neurons in their expected location, while TdTomato-expressing cells were situated more laterally (Fig. 10B, GnRH, green; TdTomato, red). There was no overlap of the green and red fluorescence in any section from either male or female mice. We also performed qPCR on biopsy punched preoptic area and analyzed *Gnrh* expression. *Gnrh* expression did not differ in control and JUN-cKO male mice (Fig. 10C). However, *Gnrh* expression was reduced by 56% in the female JUN-cKO mice (Fig. 10D). Given that GnRH neurons of either sex did not express TdTomato reporter, we hypothesize that lower *Gnrh* mRNA levels in the female may stem from the upstream regulatory neurons that may be affected by either lack of JUN in GnRH receptor-expressing cells, or by lower estrogen levels.

Discussion

The molecular mechanisms of GnRH regulation of its target genes in pituitary gonadotropes have been previously examined primarily in cell lines and in primary cultures; however, a role of GnRH-induced transcription factors regulating gonadotrope genes *in vivo* is just beginning to emerge. As an immediate-early gene, JUN is rapidly induced in gonadotrope cells following GnRH treatment, both *in vivo* (59) and in model cell lines (14, 60). In these, JUN mediates GnRH induction of the FSH β (*Fshb*) gonadotropin subunit (9) and of the GnRH receptor (*Gnrhr*) (12) by binding to the AP-1 site in the proximal promoters of these genes following dimerization with FOS. Herein, we examined the role of JUN in HPG axis gene expression *in vivo*, using c-Jun^{flox/flox} mice, crossed to GRIC, in which Cre expression is driven by the GnRH receptor promoter. We demonstrate that

JUN expression in the GnRH receptor-expressing cells is necessary for normal reproductive function.

Mice lacking JUN in GnRH receptor-expressing cells exhibit a number of reproductive defects. Males have decreased *Lhb* and *Gnrhr* expression, which results in a decline in LH concentration in the circulation, and consequent reduction in testicular function, including lower expression of several steroidogenic enzymes, leading to reduced testosterone levels, smaller seminal vesicles and fewer mature spermatozoa. Females, as well, have lower LH, which results in longer estrous cycles, reduced expression of *Cyp17* steroidogenic enzyme and fewer corpora lutea in the ovaries. Reduced number of corpora lutea, despite normal numbers of antral follicles, may stem from diminished intra-ovarian steroid hormone levels due to lower expression of *Cyp17*. Alternatively, fewer corpora lutea may be a result of abrogated prolactin levels. Prolactin has a critical permissive role for LH action in the ovary, and is necessary for luteinization and corpus luteum function in rodents (49, 50). Reduced prolactin expression likely derives from decreased levels of steroid hormones. Estrogen strongly upregulates prolactin in females (61, 62). In males, expression of aromatase in the pituitary allows for testosterone conversion to estrogen, which then increases prolactin levels (63). Therefore, decreased estrogen may contribute to diminished prolactin expression and reduced number of corpora lutea. Previous studies analyzing regulation of *Lhb* expression failed to find a role for JUN, while JUN is involved in *Gnrhr* induction. Since *Lhb* expression is dependent on GnRH receptor numbers at the surface of gonadotropes (44), we believe that reduced levels of GnRH receptors are a cause of diminished LH levels. On the other hand, it is possible that AP-1 may play a role in *Lhb* expression *in vivo*.

Unexpectedly, FSH levels were unchanged in the cKO animals, although in L β T2 model cell line JUN mediates GnRH induction of the *Fshb* subunit (9). This may illustrate discrepancy

between cell models and *in vivo* function, as suggested before (64). GRIC model has been used in the recent literature to analyze a role of transcription factors in the gonadotrope (40-42). We determined a significant overlap between LH and TdTomato expression. Although difference in the percent coexpression of the reporters and LH between previously reported results (34) and results reported herein is small, it may stem from different levels of fluorescent reporter expression. 12% of LH-expressing cells lack TdTomato expression demonstrating insufficient Cre activity in these cells. The number of FSH-expressing cells which do not have sufficient Cre expression is higher, at 24%. It is possible that FSH-containing cells that do not express functional Cre are sufficient to maintain normal levels of FSH in the circulation. Especially since FSH can be constitutively secreted (65, 66) and thus, would be less dependent on the level of GnRH receptor expression. Lack of GnRH receptor expression in a portion of the FSH-containing cells was reported previously (35), although it was postulated that this population is present only during development. Data presented herein imply that FSH-containing cells without GnRH receptor persist in adulthood, which is consistent with several previous studies (67, 68). Compensation by JUNB may explain unchanged FSH levels as well, although in most tissues JUN and JUNB have opposing effects (69, 70). However, FSH β was more highly induced by JUN heterodimers than JUNB heterodimers, while GnRH receptor induction was the same with either JUN or JUNB heterodimers with FOS or FOSB. The effect on GnRH receptor expression may indicate that either the GnRH receptor is more sensitive to the levels of JUN, or that JUNB cannot compensate for JUN to induce GnRH receptor expression.

Given that GRIC allele also drives Cre expression in the testes, in the germ cells (35), there is a concern that gonadal phenotype in the male mice may be caused by a lack of JUN in testes. However, that is unlikely for several reasons. Specific lack of JUN in male and female JUN-cKO

results in the same outcomes: lower expression of *Gnrhr* and *Lhb* mRNA in the pituitary, reduced LH in the circulation and diminished expression of LH-dependent genes in the gonads, resulting in lower sex steroid levels. In fact, in the female JUN-cKO, all these effects are exacerbated compared to the male JUN-cKO. Furthermore, known targets of AP-1 in the testes are not affected. Although AP-1 binding site was identified in the FSH receptor promoter, regulating expression of the FSH receptor by FSH (71), in the testes of JUN-cKO males FSH receptor expression is not affected (data not shown). AP-1 factors also play roles in tight junction formation and blood-testis barrier (45, 46). Blood-testis barrier, which is necessary for spermatogenesis and fertility, is established via expression of tight junction proteins, primarily Claudin 11 (72-74). Expression of Claudin 11 is unaltered in JUN-cKO mice (data not shown). Claudin 3, whose expression is regulated by androgens, forms the stage-specific basal barrier in mice (47). Despite a decrease in circulating testosterone, expression of Claudin 3 is not significantly changed either. Since late stage spermatogenesis markers are reduced, AP-1 may regulate spermatogenesis directly (75). Due to a lack of known AP-1 target genes in germ cells, we are not able to delineate if decrease in late stage spermatogenesis may be due to testicular expression of Cre or to reduced levels of LH and diminished testosterone. Therefore, since males and females JUN-cKO exhibit similar phenotypes, and Cre is not expressed in the ovary, the observed effects likely stem from the gonadotrope specific JUN knockdown.

Cre expression is driven by the GnRH receptor regulatory region, which is expressed in several other extrapituitary sites. GnRH receptors, in addition to pituitary gonadotrope, are expressed in the mediobasal hypothalamus, amygdala and hippocampus (52), but the specific neuronal populations that express GnRH receptors are not known. Several studies identified that GnRH receptor is expressed in about 50% of GnRH neurons (55-58), suggesting that GnRH

receptor may contribute to autocrine GnRH pulse generation (76, 77). Using this same GRIC mouse, ablation of GnRH receptor-expressing neurons resulted in elevated number of GnRH neurons (78), implying that GnRH receptor is not expressed in GnRH neurons themselves, but it may be expressed in afferent neurons that regulate GnRH neurons. It was also postulated that central GnRH via hypothalamic GnRH receptors upstream of GnRH neurons, may participate in the pulsatile release and preovulatory surge (79). Our analyses of GnRH receptor-driven Cre expression in the hypothalamus, demonstrated Cre activity in the arcuate nucleus and in the preoptic area, but not in GnRH neurons themselves. Examination of GnRH expression determined that *Gnrh* mRNA is significantly reduced specifically in female JUN-cKO. We previously observed female specific effects using FOS null animals (25). Since there was no overlap between GnRH neurons and TdTomato expression, these findings suggest that GnRH expression is mediated in part via activity-regulated gene induction by afferent neurons which may be affected by reduced estrogen levels or by JUN knockdown. A number of previous reports determined that hypothalamic factors involved in reproductive function, such as RFamide-related peptide 3 (RFRP-3), a mammalian gonadotropin-inhibitory hormone ortholog; senktide, a neurokinin B receptor agonist; and oxytocin; elicit changes in LH serum levels, not only via alterations of GnRH secretion but by modifications of *Gnrh* transcription (80-82). Alternatively, diminished *Gnrh* mRNA transcription may be secondary to reduced LH levels that caused lower estrogen (83, 84). Similar to other studies using whole animal models where endocrine loops are dysregulated, we are not able to distinguish between these alternatives. These results may indicate that the observed reproductive phenotype in females may stem from reduced GnRH expression.

In summary, our analyses of the mice that lack JUN in GnRH receptor-expressing cells revealed several physiological roles of this gene in the reproductive axis. Reduced GnRH receptor

and lower LH levels contribute to diminished sex-steroid hormone levels, impaired spermatogenesis and reduced numbers of corpora lutea. Unchanged FSH levels may be due to compensatory role of JUNB for this gene target but not for GnRH receptor, or to the presence of FSH gonadotropes that lack sufficient Cre activity. We demonstrate that JUN expression in GnRH receptor-expressing cells is necessary for normal reproductive function.

Acknowledgements

The authors declare no competing financial interests.

This research was supported by NIH grants R01 HD057549 and UCR School of Medicine start-up funds (to DC). University of Virginia, Ligand Core is supported by U54-HD28934.

The authors thank Dr. Randy Johnson for the cJun^{flox/flox} mice. We thank Dr. Greg Anderson for GnRH antibody and Dr. Pamela Mellon for LβT2 cells. We are grateful to Dr. Ameae Walker for thoughtful discussion.

References

1. Stojilkovic SS, Krsmanovic LZ, Spergel DJ, Catt KJ. Gonadotropin-releasing hormone neurons: Intrinsic pulsatility and receptor-mediated regulation. *Trends Endo Metab* 1994; 5:201-209
2. Stojilkovic SS, Reinhart J, Catt KJ. Gonadotropin-releasing hormone receptors: Structure and signal transduction pathways. *Endocrine Rev* 1994; 15:462-499
3. Thackray VG, Mellon PL, Coss D. Hormones in synergy: Regulation of the pituitary gonadotropin genes. *Mol Cell Endocrinol* 2010; 314:192-203
4. Coss D. Regulation of reproduction via tight control of gonadotropin hormone levels. *Mol Cell Endocrinol* 2017;
5. Ortolano GA, Haisenleder DJ, Dalkin AC, Iliff-Sizemore SA, Landefeld TD, Maurer RA, Marshall JC. Follicle-stimulating hormone beta subunit messenger ribonucleic acid concentrations during the rat estrous cycle. *Endocrinology* 1988; 123:2946-2948

- 500 6. Halvorson LM, Weiss J, Bauer-Dantoin AC, Jameson JL. Dynamic regulation of pituitary
501 follistatin messenger ribonucleic acids during the rat estrous cycle. *Endocrinology* 1994;
502 134:1247-1253
- 503 7. Pierce JG, Parsons TF. Glycoprotein hormones: structure and function. *Ann Rev*
504 *Biochem* 1981; 50:465-495
- 505 8. Karin M, Liu Z, Zandi E. AP-1 function and regulation. *Curr Opin Cell Biol* 1997; 9:240-
506 246
- 507 9. Coss D, Jacobs SB, Bender CE, Mellon PL. A novel AP-1 site is critical for maximal
508 induction of the follicle-stimulating hormone beta gene by gonadotropin-releasing
509 hormone. *J Biol Chem* 2004; 279:152-162
- 510 10. Wang Y, Fortin J, Lamba P, Bonomi M, Persani L, Roberson MS, Bernard DJ. Activator
511 protein-1 and smad proteins synergistically regulate human follicle-stimulating hormone
512 beta-promoter activity. *Endocrinology* 2008; 149:5577-5591
- 513 11. Duval DL, Nelson SE, Clay CM. The tripartite basal enhancer of the gonadotropin-
514 releasing hormone (GnRH) receptor gene promoter regulates cell-specific expression
515 through a novel GnRH receptor activating sequence. *Mol Endocrinol* 1997; 11:1814-
516 1821
- 517 12. White BR, Duval DL, Mulvaney JM, Roberson MS, Clay CM. Homologous regulation of
518 the gonadotropin-releasing hormone receptor gene is partially mediated by protein kinase
519 C activation of an Activator Protein-1 element. *Mol Endocrinol* 1999; 13:566-577
- 520 13. Norwitz ER, Xu S, Xu J, Spiryda LB, Park JS, Jeong KH, McGee EA, Kaiser UB. Direct
521 binding of AP-1 (Fos/Jun) proteins to a SMAD binding element facilitates both GnRH-
522 and activin-mediated transcriptional activation of the mouse GnRH receptor gene. *J Biol*
523 *Chem* 2002; 277:37469-37478
- 524 14. Wurmbach E, Yuen T, Ebersole BJ, Sealfon SC. Gonadotropin-releasing hormone
525 receptor-coupled gene network organization. *The Journal of biological chemistry* 2001;
526 276:47195-47201
- 527 15. Yuen T, Choi SG, Pincas H, Waring DW, Sealfon SC, Turgeon JL. Optimized
528 amplification and single-cell analysis identify GnRH-mediated activation of Rap1b in
529 primary rat gonadotropes. *Molecular and Cellular Endocrinology* 2012; 350:10-19
- 530 16. Lindaman LL, Yeh DM, Xie C, Breen KM, Coss D. Phosphorylation of ATF2 and
531 interaction with NFY induces c-Jun in the gonadotrope. *Mol Cell Endocrinol* 2013;
532 365:316-326
- 533 17. Ely HA, Mellon PL, Coss D. GnRH Induces the c-Fos gene via phosphorylation of SRF
534 by the calcium/calmodulin kinase II pathway. *Mol Endocrinol* 2011; 25:669-680

- 535 **18.** Reddy GR, Xie C, Lindaman LL, Coss D. GnRH increases c-Fos half-life contributing to
536 higher FSHbeta induction. *Mol Endocrinol* 2013; 27:253-265
- 537 **19.** Jonak CR, Lainez NM, Roybal LL, Williamson AD, Coss D. c-JUN Dimerization Protein
538 2 (JDP2) Is a Transcriptional Repressor of Follicle-stimulating Hormone β (FSH β) and Is
539 Required for Preventing Premature Reproductive Senescence in Female Mice. *Journal of*
540 *Biological Chemistry* 2017; 292:2646-2659
- 541 **20.** Ellsworth BS, Burns AT, Escudero KW, Duval DL, Nelson SE, Clay CM. The
542 gonadotropin releasing hormone (GnRH) receptor activating sequence (GRAS) is a
543 composite regulatory element that interacts with multiple classes of transcription factors
544 including Smads, AP-1 and a forkhead DNA binding protein. *Mol Cell Endocrinol* 2003;
545 206:93-111
- 546 **21.** Jeong KH, Chin WW, Kaiser UB. Essential role of the homeodomain for pituitary
547 homeobox 1 activation of mouse gonadotropin-releasing hormone receptor gene
548 expression through interactions with c-Jun and DNA. *Mol Cell Biol* 2004; 24:6127-6139
- 549 **22.** Cheng KW, Ngan ES, Kang SK, Chow BK, Leung PC. Transcriptional down-regulation
550 of human gonadotropin-releasing hormone (GnRH) receptor gene by GnRH: role of
551 protein kinase C and activating protein 1. *Endocrinology* 2000; 141:3611-3622
- 552 **23.** Lee SL, Sadovsky Y, Swirnov AH, Polish JA, Goda P, Gavriline G, Milbrandt J.
553 Luteinizing hormone deficiency and female infertility in mice lacking the transcription
554 factor NGFI-A (Egr-1). *Science* 1996; 273:1219-1221
- 555 **24.** Topilko P, Schneider-Maunoury S, Levi G, Trembleau A, Gourdji D, Driancourt MA,
556 Rao CV, Charnay P. Multiple pituitary and ovarian defects in Krox-24 (NGFI-A, Egr-1)-
557 targeted mice. *Mol Endocrinol* 1998; 12:107-122
- 558 **25.** Xie C, Jonak CR, Kauffman AS, Coss D. Gonadotropin and kisspeptin gene expression,
559 but not GnRH, are impaired in cFOS deficient mice. *Molecular and Cellular*
560 *Endocrinology* 2015; 411:223-231
- 561 **26.** Lee WS, Smith MS, Hoffman GE. cFos Activity Identifies Recruitment of Luteinizing
562 Hormone-Releasing Hormone Neurons During the Ascending Phase of the Proestrous
563 Luteinizing Hormone Surge. *Journal of neuroendocrinology* 1992; 4:161-166
- 564 **27.** Robertson JL, Clifton DK, de la Iglesia HO, Steiner RA, Kauffman AS. Circadian
565 regulation of Kiss1 neurons: implications for timing the preovulatory gonadotropin-
566 releasing hormone/luteinizing hormone surge. *Endocrinology* 2009; 150:3664-3671
- 567 **28.** Clarkson J, d'Anglemont de Tassigny X, Moreno AS, Colledge WH, Herbison AE.
568 Kisspeptin-GPR54 signaling is essential for preovulatory gonadotropin-releasing
569 hormone neuron activation and the luteinizing hormone surge. *J Neurosci* 2008; 28:8691-
570 8697

- 571 **29.** Roybal LL, Hambarchyan A, Meadows JD, Barakat NH, Pepa PA, Breen KM, Mellon
572 PL, Coss D. Roles of Binding Elements, FOXL2 Domains, and Interactions With cJUN
573 and SMADs in Regulation of FSH β . *Molecular Endocrinology* 2014; 28:1640-1655
- 574 **30.** Corpuz PS, Lindaman LL, Mellon PL, Coss D. FoxL2 is required for activin induction of
575 the mouse and human follicle-stimulating hormone β -subunit genes. *Mol Endocrinol*
576 2010; 24:1037-1051
- 577 **31.** Xie H, Cherrington BD, Meadows JD, Witham EA, Mellon PL. Msx1 Homeodomain
578 Protein Represses the alphaGSU and GnRH Receptor Genes During Gonadotrope
579 Development. *Mol Endocrinol* 2013; 27:422-436
- 580 **32.** Johnson RS, van Lingen B, Papaioannou VE, Spiegelman BM. A null mutation at the c-
581 jun locus causes embryonic lethality and retarded cell growth in culture. *Genes Dev*
582 1993; 7:1309-1317
- 583 **33.** Li G, Gustafson-Brown C, Hanks SK, Nason K, Arbeit JM, Pogliano K, Wisdom RM,
584 Johnson RS. c-Jun is essential for organization of the epidermal leading edge. *Dev Cell*
585 2003; 4:865-877
- 586 **34.** Wen S, Schwarz JR, Niculescu D, Dinu C, Bauer CK, Hirdes W, Boehm U. Functional
587 characterization of genetically labeled gonadotropes. *Endocrinology* 2008; 149:2701-
588 2711
- 589 **35.** Wen S, Ai W, Alim Z, Boehm U. Embryonic gonadotropin-releasing hormone signaling
590 is necessary for maturation of the male reproductive axis. *Proceedings of the National*
591 *Academy of Sciences of the United States of America* 2010; 107:16372-16377
- 592 **36.** Evans MC, Rizwan M, Mayer C, Boehm U, Anderson GM. Evidence that insulin
593 signalling in gonadotrophin-releasing hormone and kisspeptin neurones does not play an
594 essential role in metabolic regulation of fertility in mice. *Journal of neuroendocrinology*
595 2014; 26:468-479
- 596 **37.** Haavisto AM, Pettersson K, Bergendahl M, Perheentupa A, Roser JF, Huhtaniemi I. A
597 supersensitive immunofluorometric assay for rat luteinizing hormone. *Endocrinology*
598 1993; 132:1687-1691
- 599 **38.** Gay VL, Midgley AR, Jr., Niswender GD. Patterns of gonadotrophin secretion associated
600 with ovulation. *Fed Proc* 1970; 29:1880-1887
- 601 **39.** Haisenleder DJ, Schoenfelder AH, Marcinko ES, Geddis LM, Marshall JC. Estimation of
602 estradiol in mouse serum samples: evaluation of commercial estradiol immunoassays.
603 *Endocrinology* 2011; 152:4443-4447
- 604 **40.** Fortin J, Boehm U, Weinstein MB, Graff JM, Bernard DJ. Follicle-stimulating hormone
605 synthesis and fertility are intact in mice lacking SMAD3 DNA binding activity and
606 SMAD2 in gonadotrope cells. *FASEB journal : official publication of the Federation of*
607 *American Societies for Experimental Biology* 2014; 28:1474-1485

- 608 **41.** Fortin J, Boehm U, Deng C-X, Treier M, Bernard DJ. Follicle-stimulating hormone
609 synthesis and fertility depend on SMAD4 and FOXL2. *The FASEB Journal* 2014;
610 28:3396-3410
- 611 **42.** Tran S, Zhou X, Lafleur C, Calderon MJ, Ellsworth BS, Kimmins S, Boehm U, Treier M,
612 Boerboom D, Bernard DJ. Impaired fertility and FSH synthesis in gonadotrope-specific
613 Foxl2 knockout mice. *Mol Endocrinol* 2013; 27:407-421
- 614 **43.** Ellsworth BS, White BR, Burns AT, Cherrington BD, Otis AM, Clay CM. c-Jun N-
615 terminal kinase activation of activator protein-1 underlies homologous regulation of the
616 gonadotropin-releasing hormone receptor gene in alpha T3-1 cells. *Endocrinology* 2003;
617 144:839-849
- 618 **44.** Ciccone NA, Kaiser UB. The biology of gonadotroph regulation. *Current opinion in*
619 *endocrinology, diabetes, and obesity* 2009; 16:321-327
- 620 **45.** Lui WY, Sze KL, Lee WM. Nectin-2 expression in testicular cells is controlled via the
621 functional cooperation between transcription factors of the Sp1, CREB, and AP-1
622 families. *J Cell Physiol* 2006; 207:144-157
- 623 **46.** Musnier A, Leon K, Morales J, Reiter E, Boulo T, Costache V, Vourc'h P, Heitzler D,
624 Oulhen N, Poupon A, Boulben S, Cormier P, Crepieux P. mRNA-selective translation
625 induced by FSH in primary Sertoli cells. *Mol Endocrinol* 2012; 26:669-680
- 626 **47.** Chakraborty P, William Buaas F, Sharma M, Smith BE, Greenlee AR, Eacker SM, Braun
627 RE. Androgen-dependent sertoli cell tight junction remodeling is mediated by multiple
628 tight junction components. *Mol Endocrinol* 2014; 28:1055-1072
- 629 **48.** Boerboom D, Kumar V, Boyer A, Wang Y, Lambrot R, Zhou X, Rico C, Boehm U,
630 Paquet M, Céleste C, Kimmins S, Bernard DJ. β -Catenin Stabilization in Gonadotropes
631 Impairs FSH Synthesis in Male Mice In Vivo. *Endocrinology* 2015; 156:323-333
- 632 **49.** Bachelot A, Carre N, Mialon O, Matelot M, Servel N, Monget P, Ahtiainen P,
633 Huhtaniemi I, Binart N. The permissive role of prolactin as a regulator of luteinizing
634 hormone action in the female mouse ovary and extragonadal tumorigenesis. *Am J Physiol*
635 *Endocrinol Metab* 2013; 305:E845-852
- 636 **50.** Stocco C. The long and short of the prolactin receptor: the corpus luteum needs them
637 both! *Biol Reprod* 2012; 86:85
- 638 **51.** Kenner L, Hoebertz A, Beil T, Keon N, Karreth F, Eferl R, Scheuch H, Szremska A,
639 Amling M, Schorpp-Kistner M, Angel P, Wagner EF. Mice lacking JunB are osteopenic
640 due to cell-autonomous osteoblast and osteoclast defects. *J Cell Biol* 2004; 164:613-623
- 641 **52.** Jennes L, Conn PM. Gonadotropin-releasing hormone and its receptors in rat brain. *Front*
642 *Neuroendocrinol* 1994; 15:51-77

- 643 **53.** Albertson AJ, Navratil A, Mignot M, Dufourny L, Cherrington B, Skinner DC.
644 Immunoreactive GnRH type I receptors in the mouse and sheep brain. *Journal of*
645 *chemical neuroanatomy* 2008; 35:326-333
- 646 **54.** Schauer C, Tong T, Petitjean H, Blum T, Peron S, Mai O, Schmitz F, Boehm U,
647 Leinders-Zufall T. Hypothalamic gonadotropin-releasing hormone (GnRH) receptor
648 neurons fire in synchrony with the female reproductive cycle. *Journal of neurophysiology*
649 2015; 114:1008-1021
- 650 **55.** Xu C, Xu XZ, Nunemaker CS, Moenter SM. Dose-dependent switch in response of
651 gonadotropin-releasing hormone (GnRH) neurons to GnRH mediated through the type I
652 GnRH receptor. *Endocrinology* 2004; 145:728-735
- 653 **56.** Todman MG, Han SK, Herbison AE. Profiling neurotransmitter receptor expression in
654 mouse gonadotropin-releasing hormone neurons using green fluorescent protein-
655 promoter transgenics and microarrays. *Neuroscience* 2005; 132:703-112
- 656 **57.** Krsmanovic LZ, Stojilkovic SS, Mertz LM, Tomic M, Catt KJ. Expression of
657 gonadotropin-releasing hormone receptors and autocrine regulation of neuropeptide
658 release in immortalized hypothalamic neurons. *Proc Natl Acad Sci USA* 1993; 90:3908-
659 3912
- 660 **58.** Van Goor F, Krsmanovic LZ, Catt KJ, Stojilkovic SS. Autocrine regulation of calcium
661 influx and gonadotropin-releasing hormone secretion in hypothalamic neurons.
662 *Biochemistry and cell biology = Biochimie et biologie cellulaire* 2000; 78:359-370
- 663 **59.** Padmanabhan V, Dalkin A, Yasin M, Haisenleder DJ, Marshall JC, Landefeld TD. Are
664 immediate early genes involved in gonadotropin-releasing hormone receptor gene
665 regulation? Characterization of changes in GnRH receptor (GnRH-R), c-fos, and c-jun
666 messenger ribonucleic acids during the ovine estrous cycle. *Biol Reprod* 1995; 53:263-
667 269
- 668 **60.** Cesnjaj M, Catt KJ, Stojilkovic SS. Coordinate actions of calcium and protein kinase-C
669 in the expression of primary response genes in pituitary gonadotrophs. *Endocrinology*
670 1994; 135:692-701
- 671 **61.** Day RN, Koike S, Sakai M, Muramatsu M, Maurer R. Both Pit-1 and the estrogen
672 receptor are required for estrogen responsiveness of the rat prolactin gene. *Mol*
673 *Endocrinol* 1990; 4:1964-1971
- 674 **62.** Ishida M, Mitsui T, Izawa M, Arita J. Activation of D2 dopamine receptors inhibits
675 estrogen response element-mediated estrogen receptor transactivation in rat pituitary
676 lactotrophs. *Mol Cell Endocrinol* 2013; 375:58-67
- 677 **63.** Garcia Barrado MJ, Blanco EJ, Carretero Hernandez M, Iglesias Osma MC, Carretero M,
678 Herrero JJ, Burks DJ, Carretero J. Local transformations of androgens into estradiol by
679 aromatase P450 is involved in the regulation of prolactin and the proliferation of pituitary
680 prolactin-positive cells. *PLoS One* 2014; 9:e101403

- 681 **64.** Huang HJ, Sebastian J, Strahl BD, Wu JC, Miller WL. Transcriptional regulation of the
682 ovine follicle-stimulating hormone-beta gene by activin and gonadotropin-releasing
683 hormone (GnRH): involvement of two proximal activator protein-1 sites for GnRH
684 stimulation. *Endocrinology* 2001; 142:2267-2274
- 685 **65.** Levine JE, Duffy MT. Simultaneous measurement of luteinizing hormone (LH)-releasing
686 hormone, LH, and follicle-stimulating hormone release in intact and short-term castrate
687 rats. *Endocrinology* 1988; 122:2211-2221
- 688 **66.** Culler MD, Negro-Vilar A. Pulsatile follicle-stimulating hormone secretion is
689 independent of luteinizing hormone-releasing hormone (LHRH): pulsatile replacement of
690 LHRH bioactivity in LHRH-immunoneutralized rats. *Endocrinology* 1987; 120:2011-
691 2021
- 692 **67.** Childs GV, Unabia G, Miller BT. Cytochemical detection of gonadotropin-releasing
693 hormone-binding sites on rat pituitary cells with luteinizing hormone, follicle-stimulating
694 hormone, and growth hormone antigens during diestrous up-regulation. *Endocrinology*
695 1994; 134:1943-1951
- 696 **68.** Childs GV, Miller BT, Miller WL. Differential effects of inhibin on gonadotropin stores
697 and gonadotropin-releasing hormone binding to pituitary cells from cycling female rats.
698 *Endocrinology* 1997; 138:1577-1584
- 699 **69.** Shaulian E, Karin M. AP-1 in cell proliferation and survival. *Oncogene* 2001; 20:2390-
700 2400
- 701 **70.** Jochum W, Passegue E, Wagner EF. AP-1 in mouse development and tumorigenesis.
702 *Oncogene* 2001; 20:2401-2412
- 703 **71.** Griswold MD, Kim JS, Tribley WA. Mechanisms involved in the homologous down-
704 regulation of transcription of the follicle-stimulating hormone receptor gene in Sertoli
705 cells. *Mol Cell Endocrinol* 2001; 173:95-107
- 706 **72.** McCabe MJ, Tarulli GA, Laven-Law G, Matthiesson KL, Meachem SJ, McLachlan RI,
707 Dinger ME, Stanton PG. Gonadotropin suppression in men leads to a reduction in
708 claudin-11 at the Sertoli cell tight junction. *Human reproduction* 2016; 31:875-886
- 709 **73.** Stanton PG. Regulation of the blood-testis barrier. *Semin Cell Dev Biol* 2016; 59:166-
710 173
- 711 **74.** Stammeler A, Lüftner BU, Kliesch S, Weidner W, Bergmann M, Middendorff R, Konrad
712 L. Highly Conserved Testicular Localization of Claudin-11 in Normal and Impaired
713 Spermatogenesis. *PLOS ONE* 2016; 11:e0160349
- 714 **75.** Araujo FC, Oliveira CA, Reis AB, Del Puerto HL, Martins AS, Reis FM. Expression of
715 the proto-oncogene c-fos and the immunolocalization of c-fos, phosphorylated c-fos and
716 estrogen receptor beta in the human testis. *Histology and histopathology* 2009; 24:1515-
717 1522

76. Witkin JW. Synchronized neuronal networks: the GnRH system. *Microsc Res Tech* 1999; 44:11-18
77. Han SK, Lee K, Bhattarai JP, Herbison AE. Gonadotrophin-releasing hormone (GnRH) exerts stimulatory effects on GnRH neurons in intact adult male and female mice. *Journal of neuroendocrinology* 2010; 22:188-195
78. Wen S, Götze IN, Mai O, Schauer C, Leinders-Zufall T, Boehm U. Genetic Identification of GnRH Receptor Neurons: A New Model for Studying Neural Circuits Underlying Reproductive Physiology in the Mouse Brain. *Endocrinology* 2011; 152:1515-1526
79. DePaolo LV, King RA, Carrillo AJ. In vivo and in vitro examination of an autoregulatory mechanism for luteinizing hormone-releasing hormone. *Endocrinology* 1987; 120:272-279
80. Grachev P, Li XF, Kinsey-Jones JS, di Domenico AL, Millar RP, Lightman SL, O'Byrne KT. Suppression of the GnRH pulse generator by neurokinin B involves a kappa-opioid receptor-dependent mechanism. *Endocrinology* 2012; 153:4894-4904
81. Salehi MS, Khazali H, Mahmoudi F, Janahmadi M. Oxytocin Intranasal Administration Affects Neural Networks Upstream of GNRH Neurons. *Journal of molecular neuroscience : MN* 2017; 62:356-362
82. Xiang W, Zhang B, Lv F, Ma Y, Chen H, Chen L, Yang F, Wang P, Chu M. The Inhibitory Effects of RFamide-Related Peptide 3 on Luteinizing Hormone Release Involves an Estradiol-Dependent Manner in Prepubertal but Not in Adult Female Mice. *Biol Reprod* 2015; 93:30
83. Thanky NR, Slater R, Herbison AE. Sex differences in estrogen-dependent transcription of gonadotropin-releasing hormone (GnRH) gene revealed in GnRH transgenic mice. *Endocrinology* 2003; 144:3351-3358
84. Dorling AA, Todman MG, Korach KS, Herbison AE. Critical role for estrogen receptor alpha in negative feedback regulation of gonadotropin-releasing hormone mRNA expression in the female mouse. *Neuroendocrinology* 2003; 78:204-209

Figure Legends

1. JUN-containing heterodimers induce both FSH β and GnRH receptor reporters in L β T2 cells. Expression vectors for AP-1 isoforms were co-transfected with the mouse FSH β (A) and GnRH receptor (B) reporters. In the separate sets of samples, cells transfected with the reporters were treated with vehicle (V) or GnRH (G, 10 nM GnRH, 5 hours). Data represents a mean of 3

independent experiments each performed in triplicate, and significant induction compared to the empty vector control is indicated with *.

2. TdTomato coexpresses with gonadotropin hormones. TdTomato reporter mice were crossed with GRIC to allow TdTomato expression in the Cre-dependent manner. Pituitaries from GRIC⁺/TdTomato⁻ mice (not shown) and from GRIC⁺/TdTomato⁺ mice, from 4 separate litters, were sectioned and stained for LH and FSH. A, 88% of LH (green) cells coexpress TdTomato (red; while arrows indicate LH cells that do not express TdTomato); B, 76% of FSH cells coexpress TdTomato. One hundred gonadotropin hormone-containing cells were counted in 3 non-overlapping fields of view in 3 sections from 3 different male and 3 female mice.

3. Fertility is profoundly affected in JUN-cKO animals. A, Representative estrous cycle changes in Cre- control females (top) and JUN-cKO female (bottom) assessed by vaginal smears for 33 days starting at 8 weeks of age (E, estrus, P, proestrus, D/M (diestrus/metestrus). B, JUN-cKO animals (black bars) have increased average cycle length (6 females per group) than Cre-controls (gray bars). C, Sperm count indicates 43% lower numbers in 8-week old JUN-cKO compared to control littermates. D, Animals were continuously present in the cages with wild-type mice of opposite sex and monitored daily for litters. JUN-cKO mice had longer time interval between litters. * indicates difference between control (Ctr, gray bars) and JUN-cKO (cKO, black bars), determined by Student's T-test followed by Tukey's HSD test.

4. JUN is not required for gonadotrope differentiation. A, Pituitaries of control (Ctr, Cre-, cJun^{flox/flox} homozygous without Cre recombinase) and JUN-cKO (cKO, Cre+, cJun^{flox/flox}

homozygous with Cre recombinase) were subjected to immunohistochemistry for LH to analyze number of gonadotrope cells. B, quantification in males and females of gonadotropes indicates that the lack of JUN has no effect on gonadotrope population (Ctr, Cre-, gray bars; cKO, Cre+, black bars).

5. Lower LH levels in JUN-cKO animals. Six 8-week old control controls (Ctr, gray bars) and six JUN-cKO littermates (cKO, black bars) were analyzed for serum gonadotropin concentration. Females were monitored for the estrous cycle stage and analyzed in diestrus. Male and female JUN-cKO mice have lower level of LH than Cre- controls (A), while FSH is unchanged (B). Consequently, sex steroid levels are lower (C). Difference (*) between control (gray bars) and JUN-cKO (black bars) were determined by Student's T-test followed by Tukey's HSD test.

6. Reduced LH β and GnRH receptor expression in JUN-cKO mice. Pituitaries from six 8-week old Cre- controls (Ctr, gray bars) and six JUN-cKO littermates (cKO, black bars) were analyzed for expression of gonadotrope genes by qPCR: *Lhb* (LH β ; A-males, B-females), *Fshb* (FSH β ; C-males, D-females), *Gnrhr* (GnRH receptor; E, F) and *Cga* (common α GSU; G, H). Statistical significance (*) between control (gray bars) and JUN-cKO (black bars) were determined by Student's T-test followed by Tukey's test.

7. Reduced seminal vesicle weight and spermatogenesis in male JUN-cKO mice. A, Seminal vesicles were dissected and measured to reveal reduced weight in 12-week old JUN-cKO males. B, Testes were homogenized and intratesticular testosterone measure. C-J, Testes were homogenized and mRNA extracted using Trizol. qPCR revealed lower expression of CYP11 (D)

and CYP17 (E) steroidogenic enzymes and lower levels of mRNA for late stage spermatogenesis markers (I, J). K-L, Histological analyses of testes following H&E stain exhibits some abnormal seminiferous tubules in JUN-cKO males. Difference (*) between control (Ctr, gray bars) and JUN-cKO (cKO, black bars) were determined by Student's T-test followed by Tukey's HSD test.

8. Reduced expression of LH target gene CYP17 and fewer corpora lutea in JUN-cKO females. A, Histological analyses of ovaries following H&E stain illustrates lower number of corpora lutea in JUN-cKO females. B, Ovaries from JUN-cKO mice were smaller and C, had fewer corpora lutea. Ovaries were homogenized and mRNA extracted using Trizol. qPCR revealed lower expression of CYP17 (D) but not CYP19 (E) steroidogenic enzymes in 12-week old JUN-cKO female mice. F, Prolactin (*Prl*) expression in the pituitary was reduced. Statistical significance (*) between control (Ctr, gray bars) and JUN-cKO (cKO, black bars) were determined by Student's T-test followed by Tukey's posthoc test.

9. Increased JUNB expression. JUNB expression in the pituitaries of the 8-week old male and female control and JUN-cKO mice was analyzed to determine if JUNB expression is elevated in compensation for the lack of JUN. * indicates statistical significance determined by Student's t-test and Tukey's posthoc analysis.

10. Cre activity in the hypothalamus. A, Coronal sections at the level of the mediobasal hypothalamus demonstrate TdTomato reporter expression (red) and Cre activity in the arcuate nucleus, and GnRH axon terminals staining in the median eminence (green). B, Coronal section of the preoptic area shows that GnRH neurons (green) do not express TdTomato reporter (red).

Male and female mice from 4 separate litters were used, and no sex differences were detected. C, GnRH expression (*Gnrh*) in the hypothalami of male mice is not altered. D, Reduced expression of *Gnrh* gene in the female JUN-cKO mice. * indicates statistical significance determined by Student's t-test and Tukey's posthoc analysis.

Table 1. Antibodies

Antibody	Species	Dilution	Provider, cat # and RRID
LH	rabbit	1:300	NHPP, AFP240580Rb; RRID:AB_2665533
FSH	rabbit	1:300	NHPP, AFP-C0972881; RRID:AB_2687903
Prolactin	rabbit	1:300	NHPP, PRL; RRID:AB_2629220
GnRH	rabbit	1:5000	Greg Anderson, Univ. of Otago; RRID:AB_2721118

Table 2. Primers

Primers	Forward	Reverse
<i>Lhb</i> (LH β)	CTGTCAACGCAACTCTGG	ACAGGAGGCAAAGCAGC
<i>Fshb</i> (FSH β)	GCCGTTTCTGCATAAGC	CAATCTTACGGTCTCGTATACC
<i>Cga</i> (aGSU)	ATTCTGGTCATGCTGTCCATGT	CAGCCCATACTGGTAGATGG
<i>Gnrhr</i> (GnRH receptor)	GCCCCCTTGCTGTACAAAGC	CCGTCTGCTAGGTAGATCATCC
<i>Prl</i> (prolactin)	TGTTCCCAGCAGTCACCAT	CAGCAACAGGAGGAGTGTC
<i>Star</i> (StAR)	GGAGGGGTGGTAGTCAGGAGA	TCCCCTGTTTCGTAGCTGCTG
<i>Cyp11</i>	AAGTATGGCCCCATTTACAGG	TGGGGTCCACGATGTAAACT
<i>Cyp17a1</i>	ATCCTTGTCACGGTGGGAGA	GGAGGTGAGTCCGGTCATTG
<i>Cyp19a1</i> (aromatase)	TTCCCATGGCAGATTCTTGTG	CGAATCGGGAGATGTAGTG
<i>Shbg</i> (ABP)	GACATTCCCCAGCCTCATGCA	TGCCTCGGAAGACAGAACCAC
<i>Cldn3</i> (claudin 3)	AACTGCGTACAAGACGAGACG	GGCACCAACGGGTATAGAAAT
<i>Gnrh</i> (GnRH)	CTACTGCTGACTGTGTGTTTG	CATCTTCTTCTGCCTGGCTTC
<i>Gapdh</i>	TGCACCACCAACTGCTTAG	GGATGCAGGGATGATGTTC





Figure 1






Figure 3




Figure 4

Figure 5




Figure 5

Figure 6

[Click here to download Figure 6.eps](#)




Figure 6




Figure 7




Figure 8

Figure 9




Figure 9




Figure 10